

## Brief report

# Glycerol as a substrate for aerobic succinate production in minimal medium with *Corynebacterium glutamicum*

Boris Litsanov, Melanie Brocker and Michael Bott\*

Institut für Bio- und Geowissenschaften, IBG-1:  
Biotechnologie, Forschungszentrum Jülich, 52425  
Jülich, Germany.

## Summary

*Corynebacterium glutamicum*, an established microbial cell factory for the biotechnological production of amino acids, was recently genetically engineered for aerobic succinate production from glucose in minimal medium. In this work, the corresponding strains were transformed with plasmid pVWEx1-*glpFKD* coding for glycerol utilization genes from *Escherichia coli*. This plasmid had previously been shown to allow growth of *C. glutamicum* with glycerol as sole carbon source. The resulting strains were tested in minimal medium for aerobic succinate production from glycerol, which is a by-product in biodiesel synthesis. The best strain BL-1/pVWEx1-*glpFKD* formed 79 mM (9.3 g l<sup>-1</sup>) succinate from 375 mM glycerol, representing 42% of the maximal theoretical yield under aerobic conditions. A specific succinate production rate of 1.55 mmol g<sup>-1</sup> (cdw) h<sup>-1</sup> and a volumetric productivity of 3.59 mM h<sup>-1</sup> were obtained, the latter value representing the highest one currently described in literature. The results demonstrate that metabolically engineered strains of *C. glutamicum* are well suited for aerobic succinate production from glycerol.

## Introduction

Glycerol is a main by-product of biodiesel and bioethanol production (Yazdani and Gonzalez, 2007). By utilizing

glycerol for the production of value-added chemicals, such as 1,3-propanediol, ethanol, amino acids or succinate, the economic efficiency of these biofuel production processes can be significantly increased (Wendisch *et al.*, 2011). Succinate is a C<sub>4</sub>-dicarboxylate offering interesting prospects for the chemical industry as feedstock for the production of a large variety of important bulk chemicals, such as tetrahydrofuran (THF), 1,4-butanediol,  $\gamma$ -butyrolactone or maleic anhydride, with a total annual production of more than 500 000 tons (Zeikus *et al.*, 1999; McKinlay *et al.*, 2007).

Bio-based succinate production from glycerol has been described for a number of natural succinate producers, such as *Basfia succiniciproducens* (Scholten and Dagele, 2008) and *Anaerobiospirillum succiniciproducens* (Lee *et al.*, 2001), as well as for metabolically engineered *Escherichia coli* strains (Blankschien *et al.*, 2010; Zhang *et al.*, 2010). The described processes are anaerobic ones (with exception of one microaerobic process) and allow a maximal succinate yield of 1 mol succinate mol<sup>-1</sup> glycerol, whereas aerobic processes can maximally yield 0.5 mol mol<sup>-1</sup>. The slow anaerobic growth with glycerol, attributed to a redox imbalance resulting from the use of intermediates for biosynthesis (Zhang *et al.*, 2010), and the low energy yield during anaerobic utilization of glycerol (Gonzalez *et al.*, 2008) result in a low volumetric productivity (0.7–1.7 mM h<sup>-1</sup>) (Blankschien *et al.*, 2010; Zhang *et al.*, 2010). This disadvantage of all described processes hinders their biotechnological implementation (Zhang *et al.*, 2010). Alternatively, aerobic succinate production from glycerol with the yeast *Yarrowia lipolytica* was recently examined (Yuzbashev *et al.*, 2010). It accumulated succinate with relatively high volumetric productivity (2.29 mM h<sup>-1</sup>) under optimal conditions (pH 7). However, complex media additives were necessary for growth and product formation which can complicate downstream processing and increase the production costs. Thus, none of the so far described succinate production processes from glycerol is perfectly suitable for industrial application.

*Corynebacterium glutamicum* is a Gram-positive soil bacterium with GRAS status (generally regarded as safe). It is used industrially for the large-scale production of

Received 21 December, 2011; revised 10 March, 2012; accepted 13 March, 2012. \*For correspondence. E-mail m.bott@fz-juelich.de; Tel. (+49) 2461 613294; Fax (+49) 2461 612710.  
doi: 10.1111/j.1751-7915.2012.00347.x

**Funding Information** We thank the Federal Ministry of Food, Agriculture, and Consumer Protection (BMELV) for financial support within the ERA-IB project 'BioProChemBB' and Volker F. Wendisch (University of Bielefeld) for providing the plasmid pVWEx1-*glpFKD*.


more than 2 million tons of L-glutamate and about 1.4 million tons of L-lysine annually (Ajinomoto Co., 2010; 2011). In addition, *C. glutamicum* strains were designed for the production of many other industrially relevant products, such as L-serine (Stolz *et al.*, 2007) and D-serine (Stäbler *et al.*, 2011), L-valine (Blombach *et al.*, 2008), 2-ketoisovalerate (Krause *et al.*, 2010), putrescine (Schneider and Wendisch, 2010), isobutanol (Smith *et al.*, 2010; Blombach *et al.*, 2011), diaminopentane (Mimitsuka *et al.*, 2007), D-lactate (Okino *et al.*, 2008a) or ethanol (Inui *et al.*, 2004). The genome of *C. glutamicum* is known (Ikeda and Nakagawa, 2003; Kalinowski *et al.*, 2003; Yukawa *et al.*, 2007) and reliable tools for genetic engineering exist (Kirchner and Tauch, 2003). Moreover, extensive knowledge on the metabolism of *C. glutamicum* is available due to 60 years of research on amino acid production (Eggeling and Bott, 2005; Burkovski, 2008). Recently metabolically engineered *C. glutamicum* strains have been described which belong to the most efficient microorganisms for succinate production from glucose either under anaerobic (Okino *et al.*, 2008b; Litsanov *et al.*, 2012a) or aerobic conditions (Litsanov *et al.*, 2012b). Under aerobic conditions *C. glutamicum* was able to synthesize succinate in minimal medium with a high specific productivity [ $1.6 \text{ mmol g (cdw)}^{-1} \text{ h}^{-1}$ ]. The strain *C. glutamicum*  $\Delta\text{sdhCAB}$  (Litsanov *et al.*, 2012b) devoid of the operon coding for succinate dehydrogenase showed relatively high succinate production but also formed large amounts of acetate as a by-product. When the genes for all known metabolic routes for acetate synthesis were deleted in the  $\Delta\text{sdhCAB}$  background, the resulting strain *C. glutamicum* BL-1 showed an 82% reduction in acetate accumulation and a 65% increase in succinate production. Further improvements were obtained by increasing  $\text{C}_3$  carboxylation and by using growth-decoupled conditions (Litsanov *et al.*, 2012b).

Naturally, *C. glutamicum* ATCC 13032 is not able to utilize glycerol as carbon source, but expression of the *E. coli* *glpFKD* operon allows the organism to grow on glycerol as sole carbon and energy source (Rittmann *et al.*, 2008). Based on our previous study on aerobic succinate production from glucose we now explored the possibility of aerobic succinate production from glycerol with *C. glutamicum* (Fig. 1).

## Results and discussion


### Utilization of glycerol for aerobic succinate production by *C. glutamicum* $\Delta\text{sdhCAB/pVWEx1-glpFKD}$

We previously showed that the strains *C. glutamicum*  $\Delta\text{sdhCAB}$  and *C. glutamicum* BL-1 are capable of aerobic succinate production from glucose (Litsanov *et al.*, 2012b). To give these strains the ability to use glycerol as carbon source, they were transformed with plasmid



**Fig. 1.** Scheme of the central metabolism of *C. glutamicum* tailored for aerobic succinate production from glycerol. Enzymes whose genes were deleted are indicated by 'X'. The reactions affected by these deletions and their products are displayed in grey. Enzymes whose genes were overexpressed are highlighted in grey boxes and the arrows for the corresponding reactions are thickened. Abbreviations: ACN, aconitase; AK, acetate kinase; CoAT, acetyl-CoA : CoA transferase; CS, citrate synthase; FUM, fumarase; GF, glycerol facilitator (from *E. coli*); GK, glycerol kinase (from *E. coli*); G-3-P DH, glycerol-3-phosphate dehydrogenase (from *E. coli*); ICD, isocitrate dehydrogenase; ICL, isocitrate lyase; MQO, malate : menaquinone oxidoreductase; MS, malate synthase; OAA, oxaloacetate; ODHC, 2-oxoglutarate dehydrogenase complex; PEP, phosphoenolpyruvate; PK, pyruvate kinase; PEPCx, PEP carboxylase; PCx, pyruvate carboxylase; PDHC, pyruvate dehydrogenase complex; PTA, phosphotransacetylase; PQO, pyruvate : menaquinone oxidoreductase; SCS, succinyl-CoA synthetase; SDH, succinate dehydrogenase.

pVWEx1-*glpFKD*, coding for a glycerol facilitator, glycerol kinase and glycerol-3-phosphate dehydrogenase from *E. coli* (Rittmann *et al.*, 2008). The resulting strains were tested for aerobic succinate production from glycerol (Fig. 1). Three independent batch cultivations with each strain in glycerol minimal medium were performed in a parallel bioreactor system. Since elevated glycerol concentrations had been shown to inhibit growth (Rittmann *et al.*, 2008), the experiments were performed with



**Fig. 2.** Biomass formation (A), substrate consumption (B), succinate production (C) and acetate production (D) of *C. glutamicum*  $\Delta$ sdhCAB/pVWEx1-glpFKD (filled squares,  $n = 3$ ) and *C. glutamicum* BL-1/pVWEx1-glpFKD (filled circles,  $n = 3$ ) on glycerol and of *C. glutamicum*  $\Delta$ sdhCAB (open squares,  $n = 4$ ) and *C. glutamicum* BL-1 (open circles,  $n = 2$ ) on glucose. The values for strains  $\Delta$ sdhCAB and BL-1 on glucose were taken from Litsanov and colleagues (2012b). The experiments were performed as aerobic batch cultivations in modified CGXII medium containing 400 mM (1200 mM carbon) glycerol or 222 mM (1332 mM carbon) glucose under constantly controlled conditions of pH 7.0 and  $pO_2 > 30\%$  using a Multifors bioreactor system as described previously (Litsanov *et al.*, 2012b). The results displayed are average data including standard deviation from at least three experiments except for *C. glutamicum* BL-1, where only mean values from two independent experiments are presented. Glucose and organic acids were quantified by HPLC as described previously. Glycerol was also analysed by HPLC and quantified by an Agilent 1100 Refractive Index Detector.

400 mM glycerol as sole carbon source. With respect to carbon (1200 mM), this concentration was slightly lower than the one (1332 mM carbon) used previously for succinate production from glucose (222 mM), but in the same range to allow a comparison. To prevent acidification of the medium by the organic acid production, the pH was kept constant at 7.0 by addition of 3 M KOH. Oxygen limitation was avoided by keeping  $pO_2$  at  $> 30\%$  saturation. During cultivation growth ( $OD_{600}$ ), glycerol consumption and organic acid production were measured and compared with data previously published for aerobic succinate production from glucose by the strains

*C. glutamicum*  $\Delta$ sdhCAB and *C. glutamicum* BL-1 (Fig. 2, Table 1). Except for the carbon source, the cultivation conditions were the same as previously published (Litsanov *et al.*, 2012b).

*Corynebacterium glutamicum*  $\Delta$ sdhCAB/pVWEx1-glpFKD grew on glycerol with a growth rate of  $0.19 \text{ h}^{-1}$  and completely consumed glycerol (369 mM) within 22 h. The specific substrate uptake rate for glycerol [ $23 \text{ mmol carbon g (cdw)}^{-1} \text{ h}^{-1}$ ] was 28% lower than the specific glucose uptake rate of strain *C. glutamicum*  $\Delta$ sdhCAB [ $32 \text{ mmol carbon g (cdw)}^{-1} \text{ h}^{-1}$ ]. The different uptake mechanisms may play a role in this context. Whereas

**Table 1.** Growth and organic acid production parameters of different *C. glutamicum* succinate producer strains during aerobic cultivation on 222 mM glucose or 400 mM glycerol as carbon source in modified CGXII medium using a Multifors bioreactor system at pH 7.0 and  $pO_2 > 30\%$  saturation.

Substrate	<i>C. glutamicum</i> strain	Growth rate ( $h^{-1}$ )	Biomass [g (cdw) $l^{-1}$ ]	Substrate consumed (mM carbon) <sup>a</sup>	Substrate uptake rate [mmol carbon g (cdw) <sup>-1</sup> $h^{-1}$ ]	Final acetate titre (mM)	Final succinate titre (mM)	Succinate production rate [mmol g <sup>-1</sup> (cdw) $h^{-1}$ ]	Yield succinate/substrate (mol mol <sup>-1</sup> )	Succinate yield (%) <sup>b</sup>
Glucose	$\Delta sdhCAB$	$0.31 \pm 0.01$	$13.1 \pm 0.5$	$1351 \pm 23$	$32 \pm 1$	$125 \pm 11$	$40 \pm 3$	$0.75 \pm 0.03$	$0.18 \pm 0.01$	$18 \pm 1$
	BL-1	0.33	13.0	1399	31	22	66	1.39	0.28	29
Glycerol	$\Delta sdhCAB/pVWEx1-glpFKD$	$0.19 \pm 0.00$	$10.0 \pm 1.3$	$1107 \pm 10$	$23 \pm 2$	$89 \pm 6$	$64 \pm 1$	$1.30 \pm 0.05$	$0.17 \pm 0.00$	$35 \pm 1$
	BL-1/pVWEx1-glpFKD	$0.22 \pm 0.02$	$10.1 \pm 2.3$	$1125 \pm 35$	$25 \pm 2$	$35 \pm 2$	$79 \pm 6$	$1.55 \pm 0.23$	$0.21 \pm 0.02$	$42 \pm 4$

**a.** Glucose concentration multiplied by factor of 6, glycerol concentration multiplied by factor of 3.

**b.** Based on the maximal theoretical succinate yields from glucose (1 mol succinate mol<sup>-1</sup> glucose) or glycerol (0.5 mol succinate mol<sup>-1</sup> glycerol) calculated for non-growing cells metabolizing the substrate exclusively via glycolysis and the oxidative TCA cycle.

All parameters describing rates were calculated for the exponential growth phase of the cultures. The data are average values including standard deviation from at least three experiments except for *C. glutamicum* BL-1 where only mean values from two independent experiments are presented. The values for strains  $\Delta sdhCAB$  and BL-1 on glucose were taken from Litsanov and colleagues (2012b).

glucose is transported actively into the cell by a high-affinity phosphotransferase system (PTS) (Mori and Shiio, 1987), glycerol enters cells of *C. glutamicum* carrying pVWEx1-*glpFKD* via facilitated diffusion mediated by the glycerol facilitator GlpF (Rittmann *et al.*, 2008). Remarkably, succinate production by strain *C. glutamicum*  $\Delta sdhCAB/pVWEx1-glpFKD$  on glycerol was significantly increased compared with that of strain *C. glutamicum*  $\Delta sdhCAB$  on glucose. A succinate titre of 64 mM (+60%) was reached with a succinate production rate of  $1.3 \text{ mmol g (cdw)}^{-1} \text{ h}^{-1}$  (+73%). The final succinate yield ( $0.17 \text{ mol mol}^{-1}$  glycerol) represented 35% of the maximal theoretical yield of  $0.5 \text{ mol succinate mol}^{-1}$  glycerol. The calculation of the maximal yield was based on resting cells which catabolized glycerol exclusively via glycolysis and the oxidative branch of the TCA cycle (Fig. 1). *Corynebacterium glutamicum*  $\Delta sdhCAB$  reached only 18% of the maximal theoretical succinate yield ( $1 \text{ mol mol}^{-1}$  glucose) during growth on glucose (Litsanov *et al.*, 2012b). This difference in the relative yields could at least partially be due to the fact that 29% less acetate was formed on glycerol (Table 1). These results suggest that the utilization of glycerol favours succinate production and lowers acetate production.

Glycerol dissimilation is initiated by its ATP-dependent phosphorylation via glycerol kinase (Durnin *et al.*, 2008; Rittmann *et al.*, 2008). In contrast, glucose is phosphorylated by the glucose-specific PTS to glucose 6-phosphate with phosphoenolpyruvate (PEP) as phosphoryl donor. High glucose uptake rates favour high pyruvate and low PEP concentrations in the cell, a situation which was shown to be unfavourable for aerobic and anaerobic succinate production by *E. coli*, because PEP is the substrate for anaplerotic  $C_3$  carboxylation via phosphoenolpyruvate carboxylase (PEPCx) or phosphoenolpyruvate carboxykinase (PEPCK) (Lin *et al.*, 2005; Zhang *et al.*, 2009). In contrast, pyruvate is the major precursor for by-product synthesis in the corresponding strains. Inactivation of the PTS system in the aerobic and anaerobic succinate producers of *E. coli* improved succinate production and decreased the accumulation of by-products (Lin *et al.*, 2005; Zhang *et al.*, 2009). *Corynebacterium glutamicum* possesses two carboxylation enzymes, phosphoenolpyruvate carboxylase (Mori and Shiio, 1985; Eikmanns *et al.*, 1989) and pyruvate carboxylase (Peters-Wendisch *et al.*, 1998). Although pyruvate carboxylase is the predominant carboxylation enzyme during aerobic growth of the wild type, PEP carboxylase also contributes to the anaplerotic flux (Petersen *et al.*, 2000). In our previous study we showed that oxaloacetate formation by PEP carboxylase becomes more important during aerobic succinate production, since overexpression of the *ppc* gene led to an increased succinate yield (Litsanov *et al.*, 2012b). Based on these observations an improved avail-



ability of PEP for PEP carboxylation and a reduced level of pyruvate might be responsible for the better succinate yield and reduced acetate formation from glycerol compared with glucose. Furthermore, the higher energy content of glycerol ( $\Delta G_r^0 = -162.84 \text{ kJ mol}^{-1} \text{ C}$ ) compared with glucose ( $\Delta G_r^0 = -152.87 \text{ kJ mol}^{-1} \text{ C}$ ) (Thauer *et al.*, 1977) due to its higher reduction status and the resulting higher ATP yield might positively influence succinate formation. The conversion of glycerol to dihydroxyacetone phosphate (DHAP) yields reducing equivalents in the form of menaquinol which can drive ATP synthesis via oxidative phosphorylation (Bott and Niebisch, 2003). In contrast, conversion of glucose to DHAP via the Embden–Meyerhof pathway is not accompanied by formation of NADH or menaquinol.

#### Utilization of glycerol for aerobic succinate production by *C. glutamicum* BL-1/pVWEx1-glpFKD

*Corynebacterium glutamicum* BL-1/pVWEx1-glpFKD, a derivative of strain  $\Delta sdhCAB$  which lacks all known genes for acetate formation (Litsanov *et al.*, 2012b), grew on glycerol with a growth rate of  $0.22 \text{ h}^{-1}$  and completely consumed 375 mM glycerol within 22 h. Absence of the genes *pqo*, *ackA*, *pta* and *cat* coding for pyruvate : menaquinone oxidoreductase (PQO), acetate kinase (AK), phosphotransacetylase (PTA) and acetyl-CoA : CoA transferase (CoAT), respectively, reduced acetate formation during growth on glycerol by 61% compared with strain  $\Delta sdhCAB/pVWEx1-glpFKD$  (Table 1). Remarkably, the final acetate concentration in strain BL-1/pVWEx1-glpFKD on glycerol was higher than that of strain BL-1 on glucose, a result which cannot be explained yet. Strain BL-1/pVWEx1-glpFKD achieved a 23% higher succinate titre ( $79 \pm 6 \text{ mM}$ ), a 19% higher specific succinate productivity [ $1.55 \text{ mmol g (cdw)}^{-1} \text{ h}^{-1}$ ] and a 24% higher yield ( $0.21 \text{ mol mol}^{-1}$ ) than strain  $\Delta sdhCAB/pVWEx1-glpFKD$ . The production results of strain BL-1/pVWEx1-glpFKD on glycerol were comparable to that of strain BL-1/pAN6-*pyc*<sup>P458S</sup>*ppc*, which represents the currently best strain of *C. glutamicum* for aerobic succinate production from glucose (Litsanov *et al.*, 2012b). Plasmid pAN6-*pyc*<sup>P458S</sup>*ppc* improves PEP/pyruvate carboxylation by combined overexpression of the genes for pyruvate carboxylase (with a P458S exchange) and PEP carboxylase.

Strain BL-1/pVWEx1-glpFKD showed a volumetric succinate productivity of  $3.59 \text{ mM h}^{-1}$ , which to our knowledge represents the highest value described in literature for aerobic succinate producers from glycerol. Additional improvements of this strain are possible by increasing anaplerotic  $C_3$  carboxylation and applying growth-limiting conditions, as shown before for aerobic succinate production from glucose (Litsanov *et al.*, 2012b). Furthermore,

the genomic integration of the *E. coli* *glpFKD* genes under control of a strong constitutive promoter such as the *tuf* promoter (Litsanov *et al.*, 2012a) could have a positive effect by eliminating the requirements for plasmid maintenance and improving strain stability. For industrial application, the *C. glutamicum* strains have to be tested and if necessary improved with respect to their tolerance to raw glycerol, which often contains growth-inhibitory impurities (Moon *et al.*, 2010).

In summary, the present study describes for the first time the efficient use of glycerol as substrate for aerobic succinate production with metabolically engineered *C. glutamicum*. The described process shows two unique features: (i) the highest known volumetric productivity of all currently described microbial strains for aerobic succinate production from glycerol and (ii) the use of minimal medium. Taken together, these two advantages and the opportunities for strain and process optimization makes the current process an interesting alternative to anaerobic fermentations.

#### Conflict of interest

None declared.

#### References

- Ajinomoto Co. (2010) Food products business [WWW document]. URL <http://www.ajinomoto.com/ir/pdf/Food-Oct2010.pdf> (retrieved 27 February 2012).
- Ajinomoto Co. (2011) Feed-use amino acids business [WWW document]. URL <http://www.ajinomoto.com/ir/pdf/Feed-useAA-Oct2011.pdf> (retrieved 27 February 2012).
- Blankschien, M.D., Clomburg, J.M., and Gonzalez, R. (2010) Metabolic engineering of *Escherichia coli* for the production of succinate from glycerol. *Metab Eng* **12**: 409–419.
- Blombach, B., Schreiner, M.E., Bartek, T., Oldiges, M., and Eikmanns, B.J. (2008) *Corynebacterium glutamicum* tailored for high-yield L-valine production. *Appl Microbiol Biotechnol* **79**: 471–479.
- Blombach, B., Riester, T., Wieschalka, S., Ziert, C., Youn, J.W., Wendisch, V.F., and Eikmanns, B.J. (2011) *Corynebacterium glutamicum* tailored for efficient isobutanol production. *Appl Environ Microbiol* **77**: 3300–3310.
- Bott, M., and Niebisch, A. (2003) The respiratory chain of *Corynebacterium glutamicum*. *J Biotechnol* **104**: 129–153.
- Burkovski, A. (ed.) (2008) *Corynebacteria: Genomics and Molecular Biology*. Norfolk, UK: Caister Academic Press.
- Durnin, G., Clomburg, J., Yeates, Z., Alvarez, P.J., Zygorakis, K., Campbell, P., and Gonzalez, R. (2008) Understanding and harnessing the microaerobic metabolism of glycerol in *Escherichia coli*. *Biotechnol Bioeng* **103**: 148–161.
- Eggeling, L., and Bott, M. (2005) *Handbook of Corynebacterium glutamicum*. Boca Raton, Florida, USA: CRC Press, Taylor & Francis Group.
- Eikmanns, B.J., Follettie, M.T., Griot, M.U., and Sinskey, A.J. (1989) The phosphoenolpyruvate carboxylase gene of

- Corynebacterium glutamicum*: molecular cloning, nucleotide sequence, and expression. *Mol Gen Genet* **218**: 330–339.
- Gonzalez, R., Murarka, A., Dharmadi, Y., and Yazdani, S.S. (2008) A new model for the anaerobic fermentation of glycerol in enteric bacteria: trunk and auxiliary pathways in *Escherichia coli*. *Metab Eng* **10**: 234–245.
- Ikedo, M., and Nakagawa, S. (2003) The *Corynebacterium glutamicum* genome: features and impacts on biotechnological processes. *Appl Microbiol Biotechnol* **62**: 99–109.
- Inui, M., Kawaguchi, H., Murakami, S., Vertes, A.A., and Yukawa, H. (2004) Metabolic engineering of *Corynebacterium glutamicum* for fuel ethanol production under oxygen-deprivation conditions. *J Mol Microbiol Biotechnol* **8**: 243–254.
- Kalinowski, J., Bathe, B., Bartels, D., Bischoff, N., Bott, M., Burkowski, A., et al. (2003) The complete *Corynebacterium glutamicum* ATCC 13032 genome sequence and its impact on the production of L-aspartate-derived amino acids and vitamins. *J Biotechnol* **104**: 5–25.
- Kirchner, O., and Tauch, A. (2003) Tools for genetic engineering in the amino acid-producing bacterium *Corynebacterium glutamicum*. *J Biotechnol* **104**: 287–299.
- Krause, F.S., Blombach, B., and Eikmanns, B.J. (2010) Metabolic engineering of *Corynebacterium glutamicum* for 2-ketoisovalerate production. *Appl Environ Microbiol* **76**: 8053–8061.
- Lee, P.C., Lee, W.G., Lee, S.Y., and Chang, H.N. (2001) Succinic acid production with reduced by-product formation in the fermentation of *Anaerobiospirillum succiniciproducens* using glycerol as a carbon source. *Biotechnol Bioeng* **72**: 41–48.
- Lin, H., Bennett, G.N., and San, K.Y. (2005) Metabolic engineering of aerobic succinate production systems in *Escherichia coli* to improve process productivity and achieve the maximum theoretical succinate yield. *Metab Eng* **7**: 116–127.
- Litsanov, B., Brocker, M., and Bott, M. (2012a) Towards homosuccinate fermentation: metabolic engineering of *Corynebacterium glutamicum* for anaerobic production of succinate from glucose and formate. *Appl Environ Microbiol* **78**: 3325–3337.
- Litsanov, B., Kabus, M., Brocker, M., and Bott, M. (2012b) Efficient aerobic succinate production from glucose in minimal medium with *Corynebacterium glutamicum*. *Microb Biotechnol* **5**: 116–128.
- McKinlay, J.B., Vieille, C., and Zeikus, J.G. (2007) Prospects for a bio-based succinate industry. *Appl Microbiol Biotechnol* **76**: 727–740.
- Mimitsuka, T., Sawai, H., Hatsu, M., and Yamada, K. (2007) Metabolic engineering of *Corynebacterium glutamicum* for cadaverine fermentation. *Biosci Biotechnol Biochem* **71**: 2130–2135.
- Moon, C., Ahn, J.H., Kim, S.W., Sang, B.I., and Um, Y. (2010) Effect of biodiesel-derived raw glycerol on 1,3-propanediol production by different microorganisms. *Appl Biochem Biotechnol* **161**: 502–510.
- Mori, M., and Shiio, I. (1985) Purification and some properties of phosphoenolpyruvate carboxylase from *Brevibacterium flavum* and its aspartate-overproducing mutant. *J Biochem* **97**: 1119–1128.
- Mori, M., and Shiio, I. (1987) Phosphoenol-pyruvate : sugar phosphotransferase systems and sugar metabolism in *Brevibacterium flavum*. *Agric Biol Chem* **51**: 2671–2678.
- Okino, S., Suda, M., Fujikura, K., Inui, M., and Yukawa, H. (2008a) Production of D-lactic acid by *Corynebacterium glutamicum* under oxygen deprivation. *Appl Microbiol Biotechnol* **78**: 449–454.
- Okino, S., Noburyu, R., Suda, M., Jojima, T., Inui, M., and Yukawa, H. (2008b) An efficient succinic acid production process in a metabolically engineered *Corynebacterium glutamicum* strain. *Appl Microbiol Biotechnol* **81**: 459–464.
- Petersen, S., de Graaf, A.A., Eggeling, L., Mollney, M., Wiechert, W., and Sahm, H. (2000) *In vivo* quantification of parallel and bidirectional fluxes in the anaplerosis of *Corynebacterium glutamicum*. *J Biol Chem* **275**: 35932–35941.
- Peters-Wendisch, P.G., Kreutzer, C., Kalinowski, J., Patek, M., Sahm, H., and Eikmanns, B.J. (1998) Pyruvate carboxylase from *Corynebacterium glutamicum*: characterization, expression and inactivation of the *pyc* gene. *Microbiology* **144**: 915–927.
- Rittmann, D., Lindner, S.N., and Wendisch, V.F. (2008) Engineering of a glycerol utilization pathway for amino acid production by *Corynebacterium glutamicum*. *Appl Environ Microbiol* **74**: 6216–6222.
- Stäbler, N., Oikawa, T., Bott, M., and Eggeling, L. (2011) *Corynebacterium glutamicum* as a host for synthesis and export of D-amino acids. *J Bacteriol* **193**: 1702–1709.
- Schneider, J., and Wendisch, V.F. (2010) Putrescine production by engineered *Corynebacterium glutamicum*. *Appl Microbiol Biotechnol* **88**: 859–868.
- Scholten, E., and Dagele, D. (2008) Succinic acid production by a newly isolated bacterium. *Biotechnol Lett* **30**: 2143–2146.
- Smith, K.M., Cho, K.-M., and Liao, J.C. (2010) Engineering *Corynebacterium glutamicum* for isobutanol production. *Appl Microbiol Biotechnol* **87**: 1045–1055.
- Stolz, M., Peters-Wendisch, P., Etterich, H., Gerharz, T., Faurie, R., Sahm, H., et al. (2007) Reduced folate supply as a key to enhanced L-serine production by *Corynebacterium glutamicum*. *Appl Environ Microbiol* **73**: 750–755.
- Thauer, R.K., Jungermann, K., and Decker, K. (1977) Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol Rev* **41**: 100–180.
- Wendisch, V.F., Lindner, S.N., and Meiswinkel, T.M. (2011) Use of glycerol in biotechnological applications. In *Biodiesel – Quality, Emissions and By-products*. Montero, G., and Stoytcheva, M. (eds). Rijeka, Croatia: InTech, pp. 305–341.
- Yazdani, S.S., and Gonzalez, R. (2007) Anaerobic fermentation of glycerol: a path to economic viability for the biofuels industry. *Curr Opin Biotechnol* **18**: 213–219.
- Yukawa, H., Omumasaba, C.A., Nonaka, H., Kos, P., Okai, N., Suzuki, N., et al. (2007) Comparative analysis of the *Corynebacterium glutamicum* group and complete genome sequence of strain R. *Microbiology* **153**: 1042–1058.
- Yuzbashev, T.V., Yuzbasheva, E.Y., Sobolevskaya, T.I., Laptev, I.A., Vybornaya, T.V., Larina, A.S., et al. (2010) Production of succinic acid at low pH by a recombinant strain of the aerobic yeast *Yarrowia lipolytica*. *Biotechnol Bioeng* **107**: 673–682.

- Zeikus, J.G., Jain, M.K., and Elankovan, P. (1999) Biotechnology of succinic acid production and markets for derived industrial products. *Appl Environ Microbiol* **51**: 545–552.
- Zhang, X., Jantama, K., Shanmugam, K.T., and Ingram, L.O. (2009) Reengineering *Escherichia coli* for succinate production in mineral salts medium. *Appl Environ Microbiol* **75**: 7807–7813.
- Zhang, X., Shanmugam, K.T., and Ingram, L.O. (2010) Fermentation of glycerol to succinate by metabolically engineered strains of *Escherichia coli*. *Appl Environ Microbiol* **76**: 2397–2401.